

FLEXADRIN CREAM



HELPS RELIEVE PAIN AND SUPPORTS RANGE OF MOTION WITHIN MINUTES.

SCHEDULING STATUS:

Not Scheduled

PROPRIETY NAME (AND DOSAGE FORM):

Flexadrin™ Cream

COMPOSITION:

Actives Ingredients:	Per 100 g
Celadrin® 250 mg	7.7 g
Menthol	2 g
Vitamin E	1.2 g
Lechithin / grapeseed	1.5 g

Inactives: water aqua, glycerine, glyceryl stearate, mentha arvensis, nipaquard PDU, treithanolamine, carbomer.

PHARMACOLOGICAL CLASSIFICATION:

Health Supplements D32.16 (Other)

DESCIPLINE OF MEDICINE:

Topical pain relief cream.

PHARMACOLOGICAL ACTION:

Celadrin® supports rapid joint cushioning, quickly alleviating inflammation, building cartilage and restoring the entire joint area.

Flexadrin™ (Celadrin®) cream study, conducted at the University of Connecticut 100% of the osteoarthritic subjects showed significant improvement in just 30 minutes and cumulative benefits throughout the remaining 30 days of the study.

INDICATIONS:

Flexadrin™ Cream is indicated as a fast acting pain relief cream that may help alleviate pains and improve the mobility of soft tissue injuries and joint flexibility. It may help the body to prevent long term wear and tear on muscles and joints.

CONTRA-INDICATIONS:

Flexadrin™ Cream is contra-indicated to any hypersensitivity to any of the ingredients mentioned in composition.

WARNINGS AND SPECIAL PRECAUTIONS:

See CONTRA-INDICATIONS

Do not apply to irritated or broken skin.

For external use only.

Do not bandage.

PREGNANCY AND LACTATION:

Not recommended during pregnancy or lactation.

DOSAGE AND DIRECTION FOR USE:

Apply 3 – 4 times daily to the affected areas.

SIDE EFFECTS:

Frequent side effects:

No known or reported frequent side effects on Flexadrin™ Cream.

Less frequent side effects:

Although not frequent some people may develop a skin rash which is due to allergic reactions to any of the ingredients mentioned in the composition.

KNOWN SYMPTOMS OF OVERDOSE AND PARTICULARS ITS TREATMENT:

If you should experience any general symptoms of allergic reaction, discontinue use.

IDENTIFICATION:

White Cream

PRESENTATION:

White Cream packaged in 100 ml white tube in an outer carton.

STORAGE INSTRUCTIONS:

Store below or at 25 °C, away from sunlight

KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER:

Will be awarded with registration by the Medicines Control Council.

NAME AND BUSINESS ADDRESS OF APPLICANT:

Kenza Health (Pty) Ltd

51 APD Office Park, Kelvin Street, Kya Sands, South Africa 2162

Customer care: 0860 103 359

www.flexadrin.co.za

DATE OF PUBLICATION OF THIS PACKAGE INSERT:

June 2017

DISCLAIMER:

This medicine has not been evaluated by the Medicines Control Council. This medicine is not intended to diagnose, treat, cure or prevent any disease.

FLEXADRIN CREAM



HELP OM PYN TE VERLIG EN ONDERSTEUN DIE OMVANG VAN BEWEGING BINNE MINUTE

BYLAE STATUS:

Nie geskeduleer nie

EIENDOMSNAAM (EN DOSISVORM):

Flexadrin Cream

SAMESTELLING:

Aktiewe Bestanddele:	Per 100 g
Celadrin® 250 mg	7.7 g
Menthol	2 g
Vitamin E	1.2 g
Lechithin / grapeseed	1.5 g

Inaktiewe: water aqua, gliserien, gliserylstearaat, mentha arvensis, nipaquard PDU, treitanolamien, karbomeer.

FARMAKOLOGIESE KLASSIFIKASIE:

Gesondheidsaanvullings D32.16 (Ander)

MEDISYNE DESKIPLINE:

Aktuele pynverligtingsroom.

FARMAKOLOGIESE WERKING:

Celadrin ® help om ontsterkte gewing te herstel, verlig inflammasie, vinnig bou kraakbeen en herstel die hele gesamentlike area.

Flexadrin™ (Celadrin®) - roomstudie, uitgevoer aan die Universiteit van Connecticut, het 100% van die osteoartritiese onderwerpe beduidend verbeter in net 30 minute en kumulatiewe voordele gedurende die oorblywende 30 dae van die studie.

INDIKASIES:

Flexadrin™ Cream word aangedui as 'n vinnige waarnemende pynverligtingsroom wat kan help om pyn te verlig en die mobiliteit van sagteweefselbeserings en gesamentlike buigsaamheid te verbeter. Dit kan die liggaam help om langtermyn slytasie op spiere en gewigte te voorkom.

KONTRA-INDIKASIES:

Flexadrin™ Cream is teen enige hipersensitiwiteit aangedui vir enige van die bestanddele wat in samestelling genoem word.

WAARSKUWINGS EN SPESIALE VOORSORGMAATREËLS:

Sien KONTRA INDIKASIES

Moet nie toedien op geïrriteerde of gebroke vel nie.

Slegs vir eksterne gebruik.

Moenie verbind nie.

SWANGERSKAP EN LAKTASIE:

Nie aanbeveel tydens swangerskap of laktasie.

DOSIS EN GEBRUIKSAANWYSINGS:

Dien 3 tot 4 keer per dag toe op die geaffekteerde areas.

NEWE-EFFEKTE:

Gereelde newe-effekte:

Geen bekende of gerapporteerde gereelde newe-effekte op Flexadrin™ Cream.

Minder gereelde newe-effekte:

Alhoewel dit nie gereeld is nie, kan sommige mense 'n veluitslag ontwikkel as gevolg van allergiese reaksies op enige van die bestanddele wat in die samestelling genoem word.

BEKENDE SIMPTOME VAN OORDOSE EN BESONDERHEDE SY TREDJE:

As u enige algemene simptome van allergiese reaksie ervaar, moet u gebruik stop.

IDENTIFIKASIE:

Wit room

VOORLEGGING:

Wit Room verpak in 100 ml wit buis in 'n buitenste karton.

BERGING INSTRUKSIES:

Bêre onder of by 25 °C, weg van sonlig.

HOU BUITE BEREIK VAN KINDERS.

REGISTRASIE NOMMER:

Sal toegeken word met registrasie deur die Medisynebeheerraad.

NAAM EN BESIGHEIDSADRES VAN AANSOEKER:

Kenza Health (Edms) Bpk

51 APD Kantoorpark, Kelvinstraat, Kya Sands, Suid-Afrika 2162

Kliëntesorg: 0860 103 359

www.flexadrin.co.za

DATUM VAN PUBLIKASIE VAN HIERDIE PAKKET INSERT:

Junie 2017

VRYWARING:

Hierdie medisyne is nie deur die Medisynebeheerraad geëvalueer nie. Hierdie medisyne is nie bedoel om enige siekte te diagnoseer, behandel, genees of voorkom nie.